

5070 1600
Attorney's Docket No. 3477.910500
PATENT 1645

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: Jeffrey L. Wrana
Serial No.: 09/744,167
Filed: January 19, 2001

For: SARA PROTEINS



Date: July 25, 2002

Commissioner for Patents
Washington, DC 20231

RECEIVED

SEP 30 2002

TECH CENTER 1600/2900

INFORMATION DISCLOSURE STATEMENT

Sir:

Attached is a list of documents on form PTO-1449, together with a copy of each identified document. It is requested that these documents be considered by the Examiner and officially made of record in accordance with 37 C.F.R. § 1.97(b) and Section 609 of the MPEP. The Commissioner is hereby authorized to charge any fee, or credit any refund, to our Deposit Account No. 50-0220.

Respectfully submitted,

Karen A. Magri
Karen A. Magri
Registration No. 41,965

Customer Number:



20792

PATENT TRADEMARK OFFICE

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner for Patents, Washington, DC 20231, on July 25, 2002.

Sloan Hobbs
Sloan Hobbs


FORM PTO-1449 U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket Number
3477.91

Serial No.
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED

SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date
January 19, 2001

Group

U. S. PATENT DOCUMENTS

Examiner Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate

FOREIGN PATENT DOCUMENTS

Document Number	Date	Country	Class	Subclass	Translation Yes No

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)

1.	Abdollah, S., Macias-Silva, M., Tsukazaki, T., Hayashi, H., Attisano, L., and Wrana, J.L. (1997). TBRI phosphorylation of Smad2 on Ser 465 and 467 is required for Smad2/Smad4 complex formation and signaling. <i>J. Biol. Chem.</i> 272, 27678-27685.
2.	Attisano, L. and Wrana, J.L. (1998). Mads and Smads in TGFβ signaling. <i>Curr. Op. Cell Biol.</i> 10, 188-194.
3.	B. Meckelein et al., (1998), <i>Mol. Brain Research</i> , v. 55, pp. 181-197.
4.	Burd et al., (1998), <i>Mol. Cell.</i> , 2, 157-162.

EXAMINER

*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.


FORM PTO-1449 U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket Number
3477.91

Serial No.
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED
SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date
January 19, 2001

Group

5.	Chen, Y., Bhushan, A., and Vale, W. (1997b). Smad8 mediates the signaling of the receptor serine kinase. <i>Proc. Natl. Acad. Sci. USA</i> 94, 12938-12943.	
6.	Chen, Y., Lebrun, J.-J., and Vale, W. (1996). Regulation of transforming growth factor β - and activin-induced transcription by mammalian Mad proteins. <i>Proc. Natl. Acad. Sci. USA</i> 93, 12992-12997.	
7.	Dennler, S., Itoh, S., Vivien, D., ten Dijke, P., Huet, S., and Gauthier, J.-M. (1998). Direct binding of Smad3 and Smad4 to critical TGF β -inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. <i>EMBO J.</i> 17, 3091-3100.	
8.	Dyson, S. and Gurdon, J.B. (1998). The Interpretation of Position in a Morphogen Gradient as Revealed by Occupancy of Activin Receptors. <i>Cell</i> 93, 557-568.	
9.	E. Labbé et al., (1998), <i>Molecular Cell</i> , v. 2, pp. 109-120.	
10.	EMBL Sequence Database Accession No. EMHUM2.AB002303, KIAA0305	
11.	EMBL Sequence Database, Accession No. EMHUM1:AF130419	
12.	Faux, M. and Scott, J.D. (1996). Molecular glue: kinase anchoring and scaffold proteins. <i>Cell</i> 85, 9-12.	
13.	Gaullier et al., (1998), <i>Nature</i> , 394, 432-433.	
14.	Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997). TGF- β signaling from cell membrane to nucleus through SMAD proteins. <i>Nature</i> . 390, 465-471.	
15.	Henis, Y.I., Moustakas, A., Lin, H.Y., and Lodish, H.F. (1994). The type II and III transforming growth factor- β receptors form homo-oligomers. <i>J. Cell Biol.</i> 126, 139-154.	
16.	Hoodless, P.A., Haerry, T., Abdollah, S., Stapleton, M., O'Connor, M.B., Attisano, L., and Wrana, J.L. (1996). MADR1, a MAD-related protein that functions in BMP2 signaling pathways. <i>Cell</i> 85, 489-500.	
17.	Kim, J., Johnson, K., Chen, H.J., Carroll, S., and Laughon, A. (1997). <i>Drosophila</i> Mad binds to DNA and directly mediates activation of <i>vestigial</i> by decapentaplegic. <i>Nature</i> 388, 304-308.	
18.	Kretzschmar, M. and Massagué, J. (1998). SMADs: mediators and regulators of TGF- β signaling. <i>Current Opinion in Genetics & Development</i> 8, 103-111.	
19.	Kretzschmar, M., Liu, F., Hata, A., Doody, J., and Massagué, J. (1997). The TGF- β family mediator Smad1 is phosphorylated directly and activated functionally by the BMP receptor kinase. <i>Genes Dev.</i> 11, 984-995.	

EXAMINER
*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.


FORM PTO-1449 U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket Number
3477.91

Serial No.
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED

SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date
January 19, 2001

Group

20.	Labbé, E., Silvestri, C., Hoodless, P.A., Wrana, J.L., and Attisano, L. (1998). Smad2 and Smad3 positively and negatively regulate TGF β -dependent transcription through the forkhead DNA binding protein, FAST2. <i>Molecular Cell</i> in press.
21.	Lagna, G., Hata, A., Hemmati-Brivanlou, A., and Massagué, J. (1996). Partnership between DPC4 and SMAD proteins in TGF- β signaling pathways. <i>Nature</i> 383, 832-836.
22.	Liu, X., Sun, Y., Constantinescu, S.N., Karam, E., Weinberg, R.A., and Lodish, H.F. (1997b). Transforming growth factor β -induced phosphorylation of Smad3 is required for growth inhibition and transcriptional induction in epithelial cells. <i>Proc. Natl. Acad. Sci. USA</i> 94, 10669-10764.
23.	M. Kretzschmar et al., (1998), <i>Current Opinion in Genetics & Development</i> , v. 8, pp. 103-111.
24.	Macías-Silva, M., Abdollah, S., Hoodless, P.A., Pirone, R., Attisano, L., and Wrana, J.L. (1996). MADR2 is a substrate of the TGF β receptor and its phosphorylation is required for nuclear accumulation and signaling. <i>Cell</i> 87, 1215-1224.
25.	Nagase et al., (1997), <i>DNA Research</i> , v. 4, pp. 141-150.
26.	Nakao, A., Imamura, T., Souchelnytskyi, S., Kawabata, M., Ishisaki, A., Oeda, E., Tamaki, K., Hanai, J.-i., Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997a). TGF- β receptor-mediated signaling through Smad2, Smad3 and Smad4. <i>EMBO J.</i> 16, 5353-5362.
27.	Nakayama, T., Snyder, M.A., Grewal, S.S., Tsuneizumi, K., Tabata, T., and Christian, J.L. (1998). <i>Xenopus</i> Smad8 acts downstream of BMP-4 to modulate its activity during vertebrate embryonic patterning. <i>Development</i> 125, 857-867
28.	Nishimura, R., Kato, Y., Chen, D., Harris, S.E., Mundy, G.R., and Yoneda, T. (1998). Smad5 and DPC4 are key molecules in mediating BMP-2-induced osteoblastic differentiation of the pluripotent mesenchymal precursor cell line C2C12. <i>J. Biol. Chem.</i> 273, 1872-1879.
29.	Patki et al., (1998), <i>Nature</i> , 394, 433-434.
30.	Patterson, G., Kowek, A., Wong, A., Liu, Y., and Ruvkun, G. (1997). The DAF-3 Smad protein antagonizes TGF- β -related receptor signaling in the <i>C. elegans</i> dauer pathway. <i>Genes and Dev.</i>
31.	Pawson, T. and Scott, J.D. (1997). Signaling through scaffold, anchoring and adaptor proteins. <i>Science</i> 278, 2075-2080.
32.	Reddy, A.H. (1998), <i>Nature Biotechnology</i> , v. 16, pp. 247-252

EXAMINER

*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.


FORM PTO-1449 U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket Number
3477.91

Serial No.
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED

SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date
January 19, 2001

Group

33.	Savage, C., Das, P., Finelli, A., Townsend, S., Sun, C., Baird, S., and Padgett, R. (1996). The <i>C. elegans sma-2, sma-3</i> and <i>sma-4</i> genes define a novel conserved family of TGF- β pathway components. <i>Proc. Natl. Acad. Sci. USA</i> 93, 790-794.
34.	Sekelsky, J.J., Newfeld, S.J., Raftery, L.A., Chartoff, E.H., and Gelbart, W.M. (1995). Genetic characterization and cloning of Mothers against dpp, a gene required for decapentaplegic function in <i>Drosophila melanogaster</i> . <i>Genetics</i> 139, 1347-1358.
35.	Simonsen et al., (1998), <i>Nature</i> , 394, 494-495.
36.	Souchelnytskyi, S., Tamaki, K., Engström, U., Wernstedt, C., ten Dijke, P., and Heldin, C.-H. (1997). Phosphorylation of Ser ⁴⁶⁵ and Ser ⁴⁶⁷ in the C Terminus of Smad2 Mediates Interaction with Smad4 and Is Required for Transforming Growth Factor- α Signaling. <i>J. Biol. Chem.</i> 272, 28107-28115.
37.	Stenmark et al., (1996), <i>J. Biol. Chem.</i> , v. 271, pp. 24048-24054.
38.	Tsukazaki, T., Chiang, T.A., Davison, A.F., Attisano, L., and Wrana, J.L. (1998). SARA, a FYVE domain protein that recruits Smad2 to the TGF β receptor. <i>Cell</i> , 95, 779-791.
39.	Wiedemann et al., (1998), <i>Nature</i> , 394, 426-427.
40.	Yingling, J.M., Datto, M.B., Wong, C., Frederick, J.P., Liberati, N.T., and Wang, X.-F. (1997). Tumour Suppressor Smad4 is a Transforming Growth Factor α -Inducible DNA Binding Protein. <i>Mol. Cell. Biol.</i> 17, 7019-7028.
41.	Zawel, L., Dai, J.L., Buckhaults, P., Zhou, S., Kinzler, K.W., Vogelstein, B., and Kern, S.E. (1998). Human Smad3 and Smad4 are sequence-specific transcription activators. <i>Mol. Cell</i> 1, 611-617.
42.	Zhang, Y., Musci, T., and Derynck, R. (1997). The tumor suppressor Smad4/DPC4 as a central mediator of Smad function. <i>Curr. Biol.</i> 7, 270-276.

EXAMINER

*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.